



## AUBURN GASTROENTEROLOGY

W. Keith Adkins, M.D.  
785 N Dean Rd Suite 400  
Office (334) 329-7822  
Fax (334) 329-7890

### Miralax-Gatorade Preparation

**To ensure a successful exam, please follow all instructions carefully.**

You **MUST** arrange a ride for the day of your exam. If you fail to arrange acceptable transportation, your procedure will need to be rescheduled. If you have diabetes, ask your physician for diet and medication instructions. If you are or may be pregnant, please discuss the risks and benefits of this procedure with your doctor. Please bring a list of all of your current medications, including any over-the-counter medications with you. **If you must cancel or reschedule your appointment, please call your doctor's office as soon as possible.**

#### USE THE PREP KIT FROM YOUR DOCTOR OR PURCHASE THE FOLLOWING SUPPLIES AT YOUR LOCAL PHARMACY:

- 4 Dulcolax laxative tablets containing 5 mg of bisacodyl each (NOT Dulcolax stool softener)
- 1 - 8.3 oz. bottle Miralax (238 grams)-may substitute 2-10oz bottles of Magnesium Citrate (NOT red)
- 64 oz. clear liquid (NOT red). Gatorade, G2, Gatorade Ice, Powerade or Powerade Zero are acceptable.

#### 7 DAYS BEFORE YOUR COLONOSCOPY:

If you take aspirin or NSAIDS such as Advil, Motrin, Celebrex or Ibuprofen, you may continue to take them as usual unless otherwise instructed by your physician. You should discuss this with your physician in advance of the procedure. Ask your doctor for specific instructions if you take a blood thinner like Plavix, Pradaxa, Clopidogrel, Coumadin, Warfarin, Effient, Prasugrel or LovenoX.

#### 3 DAYS BEFORE YOUR COLONOSCOPY:

Stop eating all nuts, seeds, popcorn, vegetables, and grains.

#### 1 DAY BEFORE YOUR COLONOSCOPY:

Begin a clear liquid diet after a light breakfast. Drink at least 8 glasses of water during the day to avoid dehydration.

**At 12 noon**, take 4 Dulcolax tablets. Mix 64 oz. liquid with 8.3 oz. Miralax and place in the refrigerator (DO NOT ADD ICE)

**At 5 pm**, drink one 8 oz. glass of the Miralax/Gatorade solution and continue drinking one 8 oz. glass every 15 minutes thereafter until the mixture is gone.

##### Clear Liquid:

- Gatorade, Pedialyte, or Powerade
- Clear broth or bouillon
- Coffee or tea (no milk or non-dairy creamer)
- Carbonated and non-carbonated soft drinks
- Kool-Aid or other fruit-flavored drinks
- Strained fruit juices (no pulp)
- Jell-O, popsicles, hard candy

##### Not Clear Liquid:

- No red or purple items of any kind
- No alcohol
- No milk or non-dairy creamers
- No noodles or vegetables in soup
- No juice with pulp
- No liquid you cannot see through

#### DAY OF YOUR COLONOSCOPY:

**Nothing to drink after midnight!** You may take all of your morning medications as usual with 4 oz. of water **up to 3-4 hours before your procedure.**

**-4 hours before**, Only if instructed by your physician, drink 10 oz. of Magnesium Citrate. –

**3 hours before**, stop drinking all clear liquids.

**You are ready for the exam if you followed all instructions and your stool is no longer formed, but clear or yellow liquid.**

\*\*You will receive a phone call from a nurse at EAMC, 1-2 days prior to your procedure. If you have not received a phone call, please call EAMC at (334) 528-2007 the day before your procedure. If you need to cancel or reschedule your procedure, please call EAMC Scheduling Dept at (334) 528-1215 option "0".

\*\*If you have any questions, please call Auburn Gastroenterology at (334) 329-7822

\*\*You will need a driver to be with you the **ENTIRE TIME** and take you home after the procedure.